

ORAL-EZE DENTAL EMERGENCY TOOTHACHE MEDICATION

COMPOSITION

Active ingredients: benzocaine 7.5% w/v, clove bud oil 19% v/v
& ethanol 18.8% v/v

USE

For the temporary relief of throbbing & persistent toothache due to a cavity

WARNING

Not to be used for a period exceeding 48 hours. Toothaches and open cavities indicate serious problems which need prompt attention by a dentist. If irritation develops, or a fever and infection develop, discontinue use and see your dentist or physician promptly. Do not use if you are allergic to eugenol (clove oil), benzocaine or caine type anaesthetics. Avoid use in pregnancy as safety not established in pregnancy.

DOSAGE & ADMINISTRATION

Rinse the tooth with water to remove any food particles from the cavity. Moisten a small piece of cotton wool with 1 or 2 drops of Oral-eze dental medication and place in cavity for approximately 1 minute. Avoid touching tissues other than the tooth cavity. Apply the dose not more than four times daily or as directed by a dentist or physician. Children 2 to 12 years of age should be supervised in the use of this product. Do not give to children under 2 years of age.

PRESENTATION

Topical solution, 5 ml

STORAGE

Store below 25°C in a dry place.

POISONS SCHEDULE

S2

Petrus Pharmaceuticals Pty Ltd, Level 3, IBM Building, 1060 Hay Street, West Perth WA 6005